

NAGS rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A18071	Rabbit	1 mg/ml	58740

Applications	WB
Reactivity	Human, Mouse
Dilution	WB 1: 500-2000
Storage	-20°C/1 year
Specificity	This antibody detects endogenous levels of NAGS at Human/Mouse
Source / Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Immunogen	Synthesized peptide derived from human NAGS
Uniprot No	Q8N159
Alternative names	
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Clonality	Polyclonal
Isotype	IgG
Conjugation	
Background	The N-acetylglutamate synthase gene encodes a mitochondrial enzyme that catalyzes the formation of N-acetylglutamate (NAG) from glutamate and acetyl coenzyme-A. NAG is a cofactor of carbamyl phosphate synthetase I (CPSI), the first enzyme of the urea cycle in mammals. This gene may regulate ureagenesis by altering NAG availability and, thereby, CPSI activity. Deficiencies in N-acetylglutamate synthase have been associated with hyperammonemia. [provided by RefSeq, Jul 2008],
Other	NAGS, NAGS
Product Images:	

Application Key:

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.