

## NDUAD rabbit pAb antibody

| Catalog No :                 | Source:   | Concentration : | Mol.Wt. (Da): |
|------------------------------|---|-----------------|---------------|
| A18161                       | Rabbit  | 1 mg/ml         |               |
| <b>Applications</b>          | WB,ELISA  |                 |               |
| <b>Reactivity</b>            | Human,Mouse   |                 |               |
| <b>Dilution</b>              | WB 1:500-2000 ELISA 1:5000-20000  |                 |               |
| <b>Storage</b>               | -20°C/1 year  |                 |               |
| <b>Specificity</b>           | NDUAD Polyclonal Antibody detects endogenous levels of protein.   |                 |               |
| <b>Source / Purification</b> | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |                 |               |
| <b>Immunogen</b>             | Synthesized peptide derived from human protein . at AA range: 30-110  |                 |               |
| <b>Uniprot No</b>            | Q9P0J0  |                 |               |
| <b>Alternative names</b>     |   |                 |               |
| <b>Form</b>                  | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  |                 |               |
| <b>Clonality</b>             | Polyclonal  |                 |               |
| <b>Isotype</b>               | IgG   |                 |               |
| <b>Conjugation</b>           |   |                 |               |
| <b>Background</b>            | <p>NADH:ubiquinone oxidoreductase subunit A13(NDUFA13) Homo sapiens</p> <p>This gene encodes a subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), which functions in the transfer of electrons from NADH to the respiratory chain. The protein is required for complex I assembly and electron transfer activity. The protein binds the signal transducers and activators of transcription 3 (STAT3) transcription factor, and can function as a tumor suppressor. The human protein purified from mitochondria migrates at approximately 16 kDa. Transcripts originating from an upstream promoter and capable of expressing a protein with a longer N-terminus have been found, but their biological validity has not been determined. [provided by RefSeq, Oct 2009],</p> |                 |               |
| <b>Other</b>                 | <p>NDUFA13 GRIM19 CDA016 CGI-39, NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 13 (Cell death regulatory protein GRIM-19) (Complex I-B16.6) (CI-B16.6) (Gene associated with retinoic and interferon-induced mortality 19 protein) (GRIM-19) (Gene associated with retinoic and IFN-induced mortality 19 protein) (NADH-ubiquinone oxidoreductase B16.6 subunit)</p>   |                 |               |

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

**Trademarks**

All product names and trademarks are the property of their respective owners.

**Regulatory Disclaimer**

For life science research only. Not for use in diagnostic procedures.

---

**Contact and Support:**

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.

---