

## Olfactory receptor 13C3 rabbit pAb antibody

| Catalog No : | Source: | Concentration : | Mol.Wt. (Da): |
|--------------|---------|-----------------|---------------|
| A18826       | Rabbit  | 1 mg/ml         | 35313         |

|                              |  |
|------------------------------|--|
| <b>Applications</b>          | WB,IF,ELISA  |
| <b>Reactivity</b>            | Human  |
| <b>Dilution</b>              | WB: 1:500 - 1:2000. IF: 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications.  |
| <b>Storage</b>               | -20°C/1 year   |
| <b>Specificity</b>           | Olfactory receptor 13C3 Polyclonal Antibody detects endogenous levels of Olfactory receptor 13C3 protein.  |
| <b>Source / Purification</b> | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Immunogen</b>             | The antiserum was produced against synthesized peptide derived from human OR13C3. AA range:201-250   |
| <b>Uniprot No</b>            | Q8NGS6   |
| <b>Alternative names</b>     | OR13C3; Olfactory receptor 13C3; Olfactory receptor OR9-8  |
| <b>Form</b>                  | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| <b>Clonality</b>             | Polyclonal   |
| <b>Isotype</b>               | IgG  |
| <b>Conjugation</b>           |  |
| <b>Background</b>            | olfactory receptor family 13 subfamily C member 3(OR13C3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008], |
| <b>Other</b>                 | OR13C3, Olfactory receptor 13C3  |

### Product Images:

**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation  
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster  
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All  
Species Expected

**Trademarks**

*All product names and trademarks are the property of their respective owners.*

**Regulatory Disclaimer**

*For life science research only. Not for use in diagnostic procedures.*

**Contact and Support:**

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.