

## p27 (phospho Ser10) rabbit pAb antibody

| Catalog No :                 | Source:  | Concentration : | Mol.Wt. (Da): |
|------------------------------|--|-----------------|---------------|
| A19248                       | Rabbit   | 1 mg/ml         | 22073         |
| <b>Applications</b>          | WB,ELISA   |                 |               |
| <b>Reactivity</b>            | Human,Mouse,Rat  |                 |               |
| <b>Dilution</b>              | WB: 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.  |                 |               |
| <b>Storage</b>               | -20°C/1 year   |                 |               |
| <b>Specificity</b>           | Phospho-p27 (S10) Polyclonal Antibody detects endogenous levels of p27 protein only when phosphorylated at S10.  |                 |               |
| <b>Source / Purification</b> | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |                 |               |
| <b>Immunogen</b>             | The antiserum was produced against synthesized peptide derived from human p27 Kip1 around the phosphorylation site of Ser10. AA range:1-50   |                 |               |
| <b>Uniprot No</b>            | P46527   |                 |               |
| <b>Alternative names</b>     | CDKN1B; KIP1; Cyclin-dependent kinase inhibitor 1B; Cyclin-dependent kinase inhibitor p27; p27Kip1   |                 |               |
| <b>Form</b>                  | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |                 |               |
| <b>Clonality</b>             | Polyclonal   |                 |               |
| <b>Isotype</b>               | IgG  |                 |               |
| <b>Conjugation</b>           |  |                 |               |
| <b>Background</b>            | cyclin dependent kinase inhibitor 1B(CDKN1B) Homo sapiens This gene encodes a cyclin-dependent kinase inhibitor, which shares a limited similarity with CDK inhibitor CDKN1A/p21. The encoded protein binds to and prevents the activation of cyclin E-CDK2 or cyclin D-CDK4 complexes, and thus controls the cell cycle progression at G1. The degradation of this protein, which is triggered by its CDK dependent phosphorylation and subsequent ubiquitination by SCF complexes, is required for the cellular transition from quiescence to the proliferative state. Mutations in this gene are associated with multiple endocrine neoplasia type IV (MEN4). [provided by RefSeq, Apr 2014], |                 |               |
| <b>Other</b>                 | CDKN1B, Cyclin-dependent kinase inhibitor 1B   |                 |               |

### Product Images:

**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation  
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster  
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All  
Species Expected

**Trademarks**

*All product names and trademarks are the property of their respective owners.*

**Regulatory Disclaimer**

*For life science research only. Not for use in diagnostic procedures.*

**Contact and Support:**

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.