

## p38 rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A19287	Rabbit	1 mg/ml	
<b>Applications</b>	WB,ELISA		
<b>Reactivity</b>	Human,Mouse,Rat,Chicken(testedbyyourcustomer )		
<b>Dilution</b>	WB: 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.		
<b>Storage</b>	-20°C/1 year		
<b>Specificity</b>	p38 Polyclonal Antibody detects endogenous levels of p38 protein.		
<b>Source / Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.		
<b>Immunogen</b>	Synthesized peptide derived from the C-terminal region of human p38.		
<b>Uniprot No</b>	Q16539		
<b>Alternative names</b>	MAPK14; CSBP; CSBP1; CSBP2; CSPB1; MXI2; SAPK2A; Mitogen-activated protein kinase 14; MAP kinase 14; MAPK 14; Cytokine suppressive anti-inflammatory drug-binding protein; CSAID-binding protein; CSBP; MAP kinase MXI2; MAX-interacting protein 2; Mitogen-activated protein kinase p38 alpha; MAP kinase p38 alpha; Stress-activated protein kinase 2a; SAPK2a		
<b>Form</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.		
<b>Clonality</b>	Polyclonal		
<b>Isotype</b>	IgG		
<b>Conjugation</b>			
<b>Background</b>	mitogen-activated protein kinase 14(MAPK14) Homo sapiens The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various environmental stresses and proinflammatory cytokines. The activation requires its phosphorylation by MAP kinase kinases (MKKs), or its autophosphorylation triggered by the interaction of MAP3K7IP1/TAB1 protein with this kinase. The substrates of this kinase include transcription regulator ATF2, MEF2C, and MAX, cell cycle regulator CDC25B, and tumor suppressor p53, which suggest the roles of this kinase in stress related transcription and cell cycle regulation, as well as in genotoxic stress response. Four alternatively spliced transcript variants of this gene encoding d		
<b>Other</b>	MAPK14, Mitogen-activated protein kinase 14		

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

**Trademarks**

All product names and trademarks are the property of their respective owners.

**Regulatory Disclaimer**

For life science research only. Not for use in diagnostic procedures.

---

**Contact and Support:**

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.

---