

## p47-phox (phospho Ser370) rabbit pAb antibody

| Catalog No :                 | Source:   | Concentration : | Mol.Wt. (Da): |
|------------------------------|---|-----------------|---------------|
| A19302                       | Rabbit  | 1 mg/ml         | 44683         |
| <b>Applications</b>          | WB,IHC,ELISA  |                 |               |
| <b>Reactivity</b>            | Human,Mouse,Rat   |                 |               |
| <b>Dilution</b>              | WB: 1:500 - 1:2000. IHC: 1:100 - 1:300. ELISA: 1:5000. Not yet tested in other applications.  |                 |               |
| <b>Storage</b>               | -20°C/1 year  |                 |               |
| <b>Specificity</b>           | Phospho-p47-phox (S370) Polyclonal Antibody detects endogenous levels of p47-phox protein only when phosphorylated at S370.   |                 |               |
| <b>Source / Purification</b> | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |                 |               |
| <b>Immunogen</b>             | The antiserum was produced against synthesized peptide derived from human p47 phox around the phosphorylation site of Ser370. AA range:341-390  |                 |               |
| <b>Uniprot No</b>            | P14598  |                 |               |
| <b>Alternative names</b>     | NCF1; NOXO2; SH3PXD1A; Neutrophil cytosol factor 1; NCF-1; 47 kDa autosomal chronic granulomatous disease protein; 47 kDa neutrophil oxidase factor; NCF-47K; Neutrophil NADPH oxidase factor 1; Nox organizer 2; Nox-organizing protein 2; SH3   |                 |               |
| <b>Form</b>                  | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |                 |               |
| <b>Clonality</b>             | Polyclonal  |                 |               |
| <b>Isotype</b>               | IgG   |                 |               |
| <b>Conjugation</b>           |   |                 |               |
| <b>Background</b>            | neutrophil cytosolic factor 1(NCF1) Homo sapiens The protein encoded by this gene is a 47 kDa cytosolic subunit of neutrophil NADPH oxidase. This oxidase is a multicomponent enzyme that is activated to produce superoxide anion. Mutations in this gene have been associated with chronic granulomatous disease. [provided by RefSeq, Jul 2008], |                 |               |
| <b>Other</b>                 | NCF1, Neutrophil cytosol factor 1   |                 |               |
| <b>Product Images:</b>       |   |                 |               |

**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

**Trademarks**

*All product names and trademarks are the property of their respective owners.*

**Regulatory Disclaimer**

*For life science research only. Not for use in diagnostic procedures.*

**Contact and Support:**

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.