

## p53 (phospho Ser6) rabbit pAb antibody

| Catalog No :                 | Source:  | Concentration : | Mol.Wt. (Da): |
|------------------------------|--|-----------------|---------------|
| A19331                       | Rabbit   | 1 mg/ml         | 43653         |
| <b>Applications</b>          | WB,IHC,IP,ELISA  |                 |               |
| <b>Reactivity</b>            | Human,Monkey   |                 |               |
| <b>Dilution</b>              | WB: 1:500 - 1:2000. IHC: 1:100 - 1:300. Immunoprecipitation: 2-5 ug:mg lysate. ELISA: 1:10000. Not yet tested in other applications.   |                 |               |
| <b>Storage</b>               | -20°C/1 year   |                 |               |
| <b>Specificity</b>           | Phospho-p53 (S6) Polyclonal Antibody detects endogenous levels of p53 protein only when phosphorylated at S6.  |                 |               |
| <b>Source / Purification</b> | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |                 |               |
| <b>Immunogen</b>             | The antiserum was produced against synthesized peptide derived from human p53 around the phosphorylation site of Ser6. AA range:1-50   |                 |               |
| <b>Uniprot No</b>            | P04637   |                 |               |
| <b>Alternative names</b>     | TP53; P53; Cellular tumor antigen p53; Antigen NY-CO-13; Phosphoprotein p53; Tumor suppressor p53  |                 |               |
| <b>Form</b>                  | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |                 |               |
| <b>Clonality</b>             | Polyclonal   |                 |               |
| <b>Isotype</b>               | IgG  |                 |               |
| <b>Conjugation</b>           |  |                 |               |
| <b>Background</b>            | <p>tumor protein p53(TP53) Homo sapiens This gene encodes a tumor suppressor protein containing transcriptional activation, DNA binding, and oligomerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancers such as Li-Fraumeni syndrome. Alternative splicing of this gene and the use of alternate promoters result in multiple transcript variants and isoforms. Additional isoforms have also been shown to result from the use of alternate translation initiation codons (PMIDs: 12032546, 20937277). [provided by RefSeq, Feb 2013],</p> |                 |               |
| <b>Other</b>                 | TP53, Cellular tumor antigen p53   |                 |               |

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

**Trademarks**

All product names and trademarks are the property of their respective owners.

**Regulatory Disclaimer**

For life science research only. Not for use in diagnostic procedures.

---

**Contact and Support:**

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.

---