

## PCDA3 rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A19544	Rabbit	1 mg/ml	
<b>Applications</b>	WB		
<b>Reactivity</b>	Human		
<b>Dilution</b>	WB 1:500-2000		
<b>Storage</b>	-20°C/1 year		
<b>Specificity</b>	This antibody detects endogenous levels of PCDA3 at Human		
<b>Source / Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.		
<b>Immunogen</b>	Synthesized peptide derived from human PCDA3		
<b>Uniprot No</b>	Q9Y5H8		
<b>Alternative names</b>	Protocadherin alpha-3 (PCDH-alpha-3)		
<b>Form</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.211% sodium azide.		
<b>Clonality</b>	Polyclonal		
<b>Isotype</b>	IgG		
<b>Conjugation</b>			
<b>Background</b>	<p>This gene is a member of the protocadherin alpha gene cluster, one of three related gene clusters tandemly linked on chromosome five that demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The alpha gene cluster is composed of 15 cadherin superfamily genes related to the mouse CNR genes and consists of 13 highly similar and 2 more distantly related coding sequences. The tandem array of 15 N-terminal exons, or variable exons, are followed by downstream C-terminal exons, or constant exons, which are shared by all genes in the cluster. The large, uninterrupted N-terminal exons each encode six cadherin ectodomains while the C-terminal exons encode the cytoplasmic domain. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins that most likely play a critical role in the establishment and function of specific cell-cell connections in the brain. Alternative splicing has been observed and additional variants have been suggested but their full-length nature has yet to be determined. [provided by RefSeq, Jul 2008],</p>		
<b>Other</b>	PCDHA3, PCDA3		

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

**Trademarks**

All product names and trademarks are the property of their respective owners.

**Regulatory Disclaimer**

For life science research only. Not for use in diagnostic procedures.

---

**Contact and Support:**

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.

---