

## S10A8 rabbit pAb antibody

| Catalog No :                 | Source:   | Concentration : | Mol.Wt. (Da): |
|------------------------------|---|-----------------|---------------|
| A21148                       | Rabbit  | 1 mg/ml         |               |
| <b>Applications</b>          | WB,ELISA  |                 |               |
| <b>Reactivity</b>            | Human,Rat,Mouse   |                 |               |
| <b>Dilution</b>              | WB 1:500-2000 ELISA 1:5000-20000  |                 |               |
| <b>Storage</b>               | -20°C/1 year  |                 |               |
| <b>Specificity</b>           | S10A8 Polyclonal Antibody detects endogenous levels of protein.   |                 |               |
| <b>Source / Purification</b> | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |                 |               |
| <b>Immunogen</b>             | Synthesized peptide derived from human protein . at AA range: 30-110  |                 |               |
| <b>Uniprot No</b>            | P05109  |                 |               |
| <b>Alternative names</b>     |   |                 |               |
| <b>Form</b>                  | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  |                 |               |
| <b>Clonality</b>             | Polyclonal  |                 |               |
| <b>Isotype</b>               | IgG   |                 |               |
| <b>Conjugation</b>           |   |                 |               |
| <b>Background</b>            | S100 calcium binding protein A8(S100A8) Homo sapiens The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21. This protein may function in the inhibition of casein kinase and as a cytokine. Altered expression of this protein is associated with the disease cystic fibrosis. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2016], |                 |               |
| <b>Other</b>                 | S100A8 CAGA CFAG MRP8, Protein S100-A8 (Calgranulin-A) (Calprotectin L1L subunit) (Cystic fibrosis antigen) (CFAG) (Leukocyte L1 complex light chain) (Migration inhibitory factor-related protein 8) (MRP-8) (p8) (S100 calcium-binding protein A8) (Urinary stone protein band A)   |                 |               |

### Product Images:

**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation  
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster  
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All  
Species Expected

**Trademarks**

*All product names and trademarks are the property of their respective owners.*

**Regulatory Disclaimer**

*For life science research only. Not for use in diagnostic procedures.*

**Contact and Support:**

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.