

VN2R1P rabbit pAb antibody

| Catalog No : | Source: | Concentration : | Mol.Wt. (Da): |
|------------------------------|---|-----------------|---------------|
| A23373 | Rabbit | 1 mg/ml | 83792 |
| Applications | WB,IF,ELISA | | |
| Reactivity | Human | | |
| Dilution | WB: 1:500 - 1:2000. IF: 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications. | | |
| Storage | -20°C/1 year | | |
| Specificity | VN2R1P Polyclonal Antibody detects endogenous levels of VN2R1P protein. | | |
| Source / Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. | | |
| Immunogen | The antiserum was produced against synthesized peptide derived from human CASRL1. AA range:400-449 | | |
| Uniprot No | Q8NGV9 | | |
| Alternative names | | | |
| Form | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. | | |
| Clonality | Polyclonal | | |
| Isotype | IgG | | |
| Conjugation | | | |
| Background | <p>The CARL-1 monoclonal antibody reacts with human TWEAK, a type II transmembrane TNF superfamily member with high identity to TNF in its extracellular portion. TWEAK transcript is expressed broadly in many adult and fetal tissues, however, the staining of human peripheral blood mononuclear cells with monoclonal antibodies shows a more restricted pattern. While freshly isolated PBMCs do not express detectable levels of TWEAK on their surface, IFN-gamma-stimulated blood monocytes rapidly upregulate TWEAK surface expression. TWEAK is expressed as membrane bound and secreted forms. Interaction of TWEAK with its counter-receptor promotes secretion of IL-8, activation of NF-kappaB, proliferation of endothelial cells, and apoptosis in a number of human cell lines. Initially, DR3 was thought to be a receptor for TWEAK, but further studies have shown that TWEAK could induce apoptosis via receptors distinct from DR3. While TWEAK exhibits overlapping signaling functions to TNF, it is generally less effective in inducing apoptosis, giving rise to its name, TNF-like weak inducer of apoptosis. For detection of human TWEAK by sandwich ELISA, a combination of purified CARL-2 for capture and biotinylated CARL-1 for detection is recommended.</p> | | |

Product Images:**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.
