

WFKN2 rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A23448	Rabbit	1 mg/ml	63360

Applications	WB
Reactivity	Human, Mouse
Dilution	WB 1: 500-2000
Storage	-20°C/1 year
Specificity	This antibody detects endogenous levels of WFKN2 at Human/Mouse
Source / Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Immunogen	Synthesized peptide derived from human WFKN2
Uniprot No	Q8TEU8
Alternative names	
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Clonality	Polyclonal
Isotype	IgG
Conjugation	
Background	The WFIKKN1 protein contains a WAP domain, follistatin domain, immunoglobulin domain, two tandem Kunitz domains, and an NTR domain. This gene encodes a WFIKKN1-related protein which has the same domain organization as the WFIKKN1 protein. The WAP-type, follistatin type, Kunitz-type, and NTR-type protease inhibitory domains may control the action of multiple types of proteases. [provided by RefSeq, Jul 2008],
Other	WFIKKN2 GASP1 WFIKKNRP UNQ9235/PRO31996, WFKN2

Product Images:

Application Key:

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.