

WWOX (phospho Tyr33) rabbit pAb antibody

Applications	WB,IHC,ELISA		
Reactivity	No :	Source:	Concentration :
		Human, Mouse, Rabbit	
Dilution A23486	WB: 1:500 - 1:2000. IHC: 1:100 - 1:300. ELISA: 1:10000. Not yet tested in other applications. 1 mg/ml 46677		
Storage	-20°C/1 year		
Specificity	Phospho-WWOX (Y33) Polyclonal Antibody detects endogenous levels of WWOX protein only when phosphorylated at Y33.		
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.		
Immunogen	The antiserum was produced against synthesized peptide derived from human WWOX around the phosphorylation site of Tyr33. AA range:18-67		
Uniprot No	Q96KM3		
Alternative names	WWOX; FOR; WOX1; WW domain-containing oxidoreductase; Fragile site FRA16D oxidoreductase		
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.		
Clonality	Polyclonal		
Isotype	IgG		
Conjugation			
Background	<p>disease:Defects in WWOX may be involved in esophageal squamous cell carcinoma (ESCC) [MIM:133239].,disease:Defects in WWOX may be involved in several cancer types. The gene spans the second most common chromosomal fragile site (FRA16D) which is frequently altered in cancers. Alteration of the expression and expression of some isoforms is associated with cancers. However, it is still unclear if alteration of WWOX is directly implicated in cancerogenesis or if it corresponds to a secondary effect.,domain:The WW 1 domain mediates interaction with TP53, and probably TP73, TFAP2C, LITAF and WBP1.,function:Probable oxidoreductase, which acts as a tumor suppressor and plays a role in apoptosis. May function synergistically with TP53/p53 to control genotoxic stress-induced cell death. May also play a role in tumor necrosis factor (TNF)-mediated cell death.,PTM:Phosphorylated upon genotoxic stress. Phosphorylation of Tyr-33 regulates interaction with TP53, TP73 and MAPK8. May also regulate proapoptotic activity.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family.,similarity:Contains 2 WW domains.,subcellular location:Partially localizes to the mitochondria. Translocates to the nucleus upon genotoxic stress or TNF stimulation (By similarity). Isoform 5 and isoform 6 may localize in the nucleus.,subunit:Interacts with TP53, TP73/p73 and MAPK8. Interacts with MAPT/TAU (By similarity). Forms a ternary complex with TP53 and MDM2. Interacts with ERBB4, LITAF and WBP1. May interact with COTE1/C1orf2 and SCOTIN.,tissue specificity:Widely expressed. Strongly expressed in testis, prostate, and ovary. Overexpressed in cancer cell lines. Isoform 5 and isoform 6 may only be expressed in tumor cell lines.,</p>		
Other	WWOX, WW domain-containing oxidoreductase; WW domain-containing oxidoreductase; Fragile site FRA16D oxidoreductase		

Product Images:**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.
