

## 17 $\beta$ -HSD4 rabbit pAb antibody

| Catalog No :                 | Source:   | Concentration : | Mol.Wt. (Da): |
|------------------------------|---|-----------------|---------------|
| A10019                       | Rabbit  | 1 mg/ml         | 79686         |
| <b>Applications</b>          | WB,IHC,ELISA  |                 |               |
| <b>Reactivity</b>            | Human,Mouse,Rat   |                 |               |
| <b>Dilution</b>              | WB: 1:500 - 1:2000. IHC: 1:100-1:300. ELISA: 1:20000. Not yet tested in other applications.   |                 |               |
| <b>Storage</b>               | -20°C/1 year  |                 |               |
| <b>Specificity</b>           | 17 $\beta$ -HSD4 Polyclonal Antibody detects endogenous levels of 17 $\beta$ -HSD4 protein.   |                 |               |
| <b>Source / Purification</b> | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |                 |               |
| <b>Immunogen</b>             | The antiserum was produced against synthesized peptide derived from the N-terminal region of human HSD17B4. AA range:41-90  |                 |               |
| <b>Uniprot No</b>            | P51659  |                 |               |
| <b>Alternative names</b>     | HSD17B4; EDH17B4; Peroxisomal multifunctional enzyme type 2; MFE-2; 17-beta-hydroxysteroid dehydrogenase 4; 17-beta-HSD 4; D-bifunctional protein; DBP; Multifunctional protein 2; MPF-2  |                 |               |
| <b>Form</b>                  | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |                 |               |
| <b>Clonality</b>             | Polyclonal  |                 |               |
| <b>Isotype</b>               | IgG   |                 |               |
| <b>Conjugation</b>           |   |                 |               |
| <b>Background</b>            | hydroxysteroid 17-beta dehydrogenase 4(HSD17B4) Homo sapiens The protein encoded by this gene is a bifunctional enzyme that is involved in the peroxisomal beta-oxidation pathway for fatty acids. It also acts as a catalyst for the formation of 3-ketoacyl-CoA intermediates from both straight-chain and 2-methyl-branched-chain fatty acids. Defects in this gene that affect the peroxisomal fatty acid beta-oxidation activity are a cause of D-bifunctional protein deficiency (DBPD). An apparent pseudogene of this gene is present on chromosome 8. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, May 2014], |                 |               |
| <b>Other</b>                 | HSD17B4, Peroxisomal multifunctional enzyme type 2; 17-beta-hydroxysteroid dehydrogenase 4; D-bifunctional protein  |                 |               |

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

**Trademarks**

All product names and trademarks are the property of their respective owners.

**Regulatory Disclaimer**

For life science research only. Not for use in diagnostic procedures.

---

**Contact and Support:**

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.

---