

A Cyclase V/VI rabbit pAb antibody

| Catalog No : | Source: | Concentration : | Mol.Wt. (Da): |
|------------------------------|---|-----------------|---------------|
| A10088 | Rabbit | 1 mg/ml | 130615 |
| Applications | IHC,IF,ELISA | | |
| Reactivity | Human,Mouse,Rat | | |
| Dilution | IHC: 1:100 - 1:300. IF: 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications. | | |
| Storage | -20°C/1 year | | |
| Specificity | A Cyclase V/VI Polyclonal Antibody detects endogenous levels of A Cyclase V/VI protein. | | |
| Source / Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. | | |
| Immunogen | The antiserum was produced against synthesized peptide derived from human ADCY5/6. AA range:931-980 | | |
| Uniprot No | O43306/O95622 | | |
| Alternative names | ADCY6; KIAA0422; Adenylate cyclase type 6; ATP pyrophosphate-lyase 6; Adenylate cyclase type VI; Adenylyl cyclase 6; Ca(2+)-inhibitable adenylyl cyclase; ADCY5; Adenylate cyclase type 5; ATP pyrophosphate-lyase 5; Adenylate cyclase type V; | | |
| Form | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. | | |
| Clonality | Polyclonal | | |
| Isotype | IgG | | |
| Conjugation | | | |
| Background | adenylate cyclase 6(ADCY6) Homo sapiens This gene encodes a member of the adenylyl cyclase family of proteins, which are required for the synthesis of cyclic AMP. All members of this family have an intracellular N-terminus, a tandem repeat of six transmembrane domains separated by a cytoplasmic loop, and a C-terminal cytoplasmic domain. The two cytoplasmic regions bind ATP and form the catalytic core of the protein. Adenylyl cyclases are important effectors of transmembrane signaling pathways and are regulated by the activity of G protein coupled receptor signaling. This protein belongs to a small subclass of adenylyl cyclase proteins that are functionally related and are inhibited by protein kinase A, calcium ions and nitric oxide. A mutation in this gene is associated with arthrogryposis multiplex congenita. [provided by RefSeq, May 2015], | | |
| Other | ADCY5/ADCY6, Adenylate cyclase type 5/6; Adenylate cyclase type 6; ATP pyrophosphate-lyase 6; Adenylate cyclase type VI; Adenylyl cyclase 6; Ca(2+)-inhibitable adenylyl cyclase; ADCY5; Adenylate cyclase type 5; ATP pyrophosphate-lyase 5; Adenylate cyclase type V; | | |

Product Images:**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.
