

AR (phospho Ser650) rabbit pAb antibody

| Catalog No : | Source: | Concentration : | Mol.Wt. (Da): |
|--------------|---------|-----------------|---------------|
| A10759 | Rabbit | 1 mg/ml | 98989 |

| | |
|------------------------------|---|
| Applications | IHC,ELISA |
| Reactivity | Human |
| Dilution | IHC: 1:100 - 1:300. ELISA: 1:40000. Not yet tested in other applications. |
| Storage | -20°C/1 year |
| Specificity | Phospho-AR (S213) Polyclonal Antibody detects endogenous levels of AR protein only when phosphorylated at S213. |
| Source / Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen | The antiserum was produced against synthesized peptide derived from human Androgen Receptor around the phosphorylation site of Ser213. AA range:186-235 |
| Uniprot No | P10275 |
| Alternative names | AR; DHTR; NR3C4; Androgen receptor; Dihydrotestosterone receptor; Nuclear receptor subfamily 3 group C member 4 |
| Form | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Clonality | Polyclonal |
| Isotype | IgG |
| Conjugation | |
| Background | androgen receptor(AR) Homo sapiens The androgen receptor gene is more than 90 kb long and codes for a protein that has 3 major functional domains: the N-terminal domain, DNA-binding domain, and androgen-binding domain. The protein functions as a steroid-hormone activated transcription factor. Upon binding the hormone ligand, the receptor dissociates from accessory proteins, translocates into the nucleus, dimerizes, and then stimulates transcription of androgen responsive genes. This gene contains 2 polymorphic trinucleotide repeat segments that encode polyglutamine and polyglycine tracts in the N-terminal transactivation domain of its protein. Expansion of the polyglutamine tract from the normal 9-34 repeats to the pathogenic 38-62 repeats causes spinal bulbar muscular atrophy (Kennedy disease). Mutations in this gene are also associated with complete androgen insensitivity (CAIS). Two alternatively spliced variants encoding distinct isoform |
| Other | AR, Androgen receptor; Dihydrotestosterone receptor; Nuclear receptor subfamily 3 group C member 4 |

Product Images:**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.
