

## CaSR rabbit pAb antibody

| Catalog No :                 | Source:  | Concentration : | Mol.Wt. (Da): |
|------------------------------|--|-----------------|---------------|
| A11656                       | Rabbit   | 1 mg/ml         | 120674        |
| <b>Applications</b>          | WB,IHC,ELISA   |                 |               |
| <b>Reactivity</b>            | Human,Mouse,Rat  |                 |               |
| <b>Dilution</b>              | WB: 1:500 - 1:2000. IHC: 1:100 - 1:300. ELISA: 1:5000. Not yet tested in other applications.   |                 |               |
| <b>Storage</b>               | -20°C/1 year   |                 |               |
| <b>Specificity</b>           | CaSR Polyclonal Antibody detects endogenous levels of CaSR protein.  |                 |               |
| <b>Source / Purification</b> | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |                 |               |
| <b>Immunogen</b>             | The antiserum was produced against synthesized peptide derived from human Calcium Sensing Receptor. AA range:854-903   |                 |               |
| <b>Uniprot No</b>            | P41180   |                 |               |
| <b>Alternative names</b>     | CASR; GPRC2A; PCAR1; Extracellular calcium-sensing receptor; CaSR; Parathyroid cell calcium-sensing receptor; PCaR1  |                 |               |
| <b>Form</b>                  | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |                 |               |
| <b>Clonality</b>             | Polyclonal   |                 |               |
| <b>Isotype</b>               | IgG  |                 |               |
| <b>Conjugation</b>           |  |                 |               |
| <b>Background</b>            | calcium sensing receptor(CASR) Homo sapiens The protein encoded by this gene is a G protein-coupled receptor that is expressed in the parathyroid hormone (PTH)-producing chief cells of the parathyroid gland, and the cells lining the kidney tubule. It senses small changes in circulating calcium concentration and couples this information to intracellular signaling pathways that modify PTH secretion or renal cation handling, thus this protein plays an essential role in maintaining mineral ion homeostasis. Mutations in this gene cause familial hypocalciuric hypercalcemia, familial, isolated hypoparathyroidism, and neonatal severe primary hyperparathyroidism. [provided by RefSeq, Jul 2008], |                 |               |
| <b>Other</b>                 | CASR, Extracellular calcium-sensing receptor   |                 |               |
| <b>Product Images:</b>       |  |                 |               |

**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation  
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster  
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All  
Species Expected

**Trademarks**

*All product names and trademarks are the property of their respective owners.*

**Regulatory Disclaimer**

*For life science research only. Not for use in diagnostic procedures.*

**Contact and Support:**

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.