

## CD299 rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A11986	Rabbit	1 mg/ml	

<b>Applications</b>	WB,IHC,ELISA
<b>Reactivity</b>	Human,Mouse,Rat
<b>Dilution</b>	WB 1:500-2000,IHC 1:500-200, ELISA 1:10000-20000
<b>Storage</b>	-20°C/1 year
<b>Specificity</b>	The antibody detects endogenous CD299
<b>Source / Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen</b>	Synthetic peptide from human protein at AA range: 271-320
<b>Uniprot No</b>	Q9H2X3
<b>Alternative names</b>	C-type lectin domain family 4 member M (CD209 antigen-like protein 1) (DC-SIGN-related protein) (DC-SIGNR) (Dendritic cell-specific ICAM-3-grabbing non-integrin 2) (DC-SIGN2) (Liver/lymph node-specific ICAM-3-grabbing non-integrin) (L-SIGN) (CD antigen CD299)
<b>Form</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Conjugation</b>	
<b>Background</b>	C-type lectin domain family 4 member M(CLEC4M) Homo sapiens This gene encodes a transmembrane receptor and is often referred to as L-SIGN because of its expression in the endothelial cells of the lymph nodes and liver. The encoded protein is involved in the innate immune system and recognizes numerous evolutionarily divergent pathogens ranging from parasites to viruses, with a large impact on public health. The protein is organized into three distinct domains: an N-terminal transmembrane domain, a tandem-repeat neck domain and C-type lectin carbohydrate recognition domain. The extracellular region consisting of the C-type lectin and neck domains has a dual function as a pathogen recognition receptor and a cell adhesion receptor by binding carbohydrate ligands on the surface of microbes and endogenous cells. The neck region is important for homo-oligomerization which allows the receptor to bind multivalent ligands with high avidity. Variations
<b>Other</b>	CLEC4M CD209L CD209L1 CD299, C-type lectin domain family 4 member M (CD209 antigen-like protein 1) (DC-SIGN-related protein) (DC-SIGNR) (Dendritic cell-specific ICAM-3-grabbing non-integrin 2) (DC-SIGN2) (Liver/lymph node-specific ICAM-3-grabbing non-integrin) (L-SIGN) (CD antigen CD299)

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation  
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster  
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All  
Species Expected

**Trademarks**

*All product names and trademarks are the property of their respective owners.*

**Regulatory Disclaimer**

*For life science research only. Not for use in diagnostic procedures.*

---

**Contact and Support:**

*To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).*

*To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.*

---