

CDK3 rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A12203	Rabbit	1 mg/ml	

Applications	WB,ELISA
Reactivity	Human,Mouse
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage	-20°C/1 year
Specificity	CDK3 Polyclonal Antibody detects endogenous levels of protein.
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Immunogen	Synthesized peptide derived from human protein . at AA range: 10-90
Uniprot No	Q00526
Alternative names	
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Clonality	Polyclonal
Isotype	IgG
Conjugation	
Background	cyclin dependent kinase 3(CDK3) Homo sapiens This gene encodes a member of the cyclin-dependent protein kinase family. The protein promotes entry into S phase, in part by activating members of the E2F family of transcription factors. The protein also associates with cyclin C and phosphorylates the retinoblastoma 1 protein to promote exit from G0. [provided by RefSeq, Jul 2008],
Other	CDK3 CDKN3, Cyclin-dependent kinase 3 (EC 2.7.11.22) (Cell division protein kinase 3)

Product Images:

Application Key:

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.