

## CFC1B rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A12325	Rabbit	1 mg/ml	

<b>Applications</b>	WB,ELISA
<b>Reactivity</b>	Human
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Storage</b>	-20°C/1 year
<b>Specificity</b>	CFC1B Polyclonal Antibody detects endogenous levels of protein.
<b>Source / Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Uniprot No</b>	P0CG36
<b>Alternative names</b>	
<b>Form</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Conjugation</b>	
<b>Background</b>	<p>disease:Defects in CFC1 are a cause of conotruncal heart malformations (CTHM) [MIM:217095]. CTHM consist of cardiac outflow tract defects, such as tetralogy of Fallot, pulmonary atresia, double-outlet right ventricle, truncus arteriosus communis, and aortic arch anomalies.,disease:Defects in CFC1 are a cause of transposition of the great arteries, dextro-looped (DTGA) [MIM:608808]. The more common form of DTGA, consists of complete inversion of the great vessels, so that the aorta incorrectly arises from the right ventricle and the pulmonary artery incorrectly arises from the left ventricle. This creates completely separate pulmonary and systemic circulatory systems, an arrangement that is incompatible with life. Patients often have atrial and/or ventricular septal defects or other types of shunting that allow some mixing between the circulations in order to support life minimally, but surgical intervention is always required.,disease:Defects in CFC1 are a cause of visceral heterotaxy (HTX2) [MIM:605376]. HTX2 is an autosomal form of visceral heterotaxy (HTX). HTX is characterized by a variable group of congenital anomalies that include complex cardiac malformations and situs inversus or situs ambiguus.,function:Involved in the correct establishment of the left-right axis. May play a role in mesoderm and/or neural patterning during gastrulation.,PTM:N-glycosylated.,similarity:Contains 1 EGF-like domain.,</p>

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation  
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster  
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All  
Species Expected

**Trademarks**

*All product names and trademarks are the property of their respective owners.*

**Regulatory Disclaimer**

*For life science research only. Not for use in diagnostic procedures.*

---

**Contact and Support:**

*To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).*

*To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.*

---