

CLAP2 rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A12530	Rabbit	1 mg/ml	
Applications	WB,ELISA		
Reactivity	Human		
Dilution	WB 1:500-2000 ELISA 1:5000-20000		
Storage	-20°C/1 year		
Specificity	CLAP2 Polyclonal Antibody detects endogenous levels of protein.		
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.		
Immunogen	Synthesized peptide derived from part region of human protein		
Uniprot No	O75122		
Alternative names			
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.		
Clonality	Polyclonal		
Isotype	IgG		
Conjugation			
Background	<p>alternative products:Additional isoforms exist,function:Microtubule plus-end tracking protein that promotes the stabilization of dynamic microtubules. Required for the polarization of the cytoplasmic microtubule arrays in migrating cells towards the leading edge of the cell. May act at the cell cortex to enhance the frequency of rescue of depolymerizing microtubules by attaching their plus-ends to cortical platforms composed of ERC1 and PHLDB2. This cortical microtubule stabilizing activity is regulated at least in part by phosphatidylinositol 3-kinase signaling. Also performs a similar stabilizing function at the kinetochore which is essential for the bipolar alignment of chromosomes on the mitotic spindle.,PTM:Phosphorylated by GSK3B. Phosphorylation by GSK3B may negatively regulate binding to microtubule lattices in lamella. Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the CLASP family.,similarity:Contains 5 HEAT repeats.,subcellular location:Localizes to microtubule plus ends. Localizes to centrosomes, kinetochores and the mitotic spindle from prometaphase. Subsequently localizes to the spindle midzone from anaphase and to the midbody from telophase. In migrating cells localizes to the plus ends of microtubules within the cell body and to the entire microtubule lattice within the lamella. Localizes to the cell cortex and this requires ERC1 and PHLDB2.,subunit:Interacts with CLIP2, ERC1, MAPRE1, MAPRE3, microtubules, PHLDB2 and RSN. The interaction with ERC1 may be mediated by PHLDB2.,tissue specificity:Brain-specific.</p>		

Product Images:**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.
