

CMTM8 rabbit pAb antibody

| Catalog No : | Source: | Concentration : | Mol.Wt. (Da): |
|--------------|---------|-----------------|---------------|
| A12727 | Rabbit | 1 mg/ml | 19572 |

| | |
|------------------------------|--|
| Applications | WB,IHC,ELISA |
| Reactivity | Human |
| Dilution | WB: 1:500 - 1:2000. IHC: 1:100-1:300. ELISA: 1:10000. Not yet tested in other applications. |
| Storage | -20°C/1 year |
| Specificity | CMTM8 Polyclonal Antibody detects endogenous levels of CMTM8 protein. |
| Source / Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen | The antiserum was produced against synthesized peptide derived from the Internal region of human CMTM8. AA range:101-150 |
| Uniprot No | Q8IZV2 |
| Alternative names | CMTM8; CKLFSF8; CKLF-like MARVEL transmembrane domain-containing protein 8; Chemokine-like factor superfamily member 8 |
| Form | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Clonality | Polyclonal |
| Isotype | IgG |
| Conjugation | |
| Background | CKLF like MARVEL transmembrane domain containing 8(CMTM8) Homo sapiens This gene belongs to the chemokine-like factor gene superfamily, a novel family that is similar to the chemokine and the transmembrane 4 superfamilies. This gene is one of several chemokine-like factor genes located in a cluster on chromosome 3. This gene acts as a tumor suppressor, and plays a role in regulating the migration of tumor cells. The encoded protein is thought to function as a negative regulator of epidermal growth factor-induced signaling. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Feb 2016], |
| Other | CMTM8, CKLF-like MARVEL transmembrane domain-containing protein 8 |
| Product Images: | |

Application Key:

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.