

## c-Myc (phospho Ser62) rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A12734	Rabbit	1 mg/ml	48804
<b>Applications</b>	WB,IHC,IF,ELISA		
<b>Reactivity</b>	Human,Mouse,Rat		
<b>Dilution</b>	WB: 1:500 - 1:2000. IHC: 1:100 - 1:300. IF: 1:200 - 1:1000. ELISA: 1:40000. Not yet tested in other applications.		
<b>Storage</b>	-20°C/1 year		
<b>Specificity</b>	Phospho-c-Myc (S62) Polyclonal Antibody detects endogenous levels of c-Myc protein only when phosphorylated at S62.		
<b>Source / Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.		
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human Myc around the phosphorylation site of Ser62. AA range:31-80		
<b>Uniprot No</b>	P01106		
<b>Alternative names</b>	MYC; BHLHE39; Myc proto-oncogene protein; Class E basic helix-loop-helix protein 39; bHLHe39; Proto-oncogene c-Myc; Transcription factor p64		
<b>Form</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.		
<b>Clonality</b>	Polyclonal		
<b>Isotype</b>	IgG		
<b>Conjugation</b>			
<b>Background</b>	v-myc avian myelocytomatosis viral oncogene homolog(MYC) Homo sapiens The protein encoded by this gene is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. There is evidence to show that alternative translation initiations from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site result in the production of two isoforms with distinct N-termini. The synthesis of non-AUG initiated protein is suppressed in Burkitt's lymphomas, suggesting its importance in the normal function of this gene. [provided by RefSeq, Jul 2008],		
<b>Other</b>	MYC, Myc proto-oncogene protein		

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

**Trademarks**

All product names and trademarks are the property of their respective owners.

**Regulatory Disclaimer**

For life science research only. Not for use in diagnostic procedures.

---

**Contact and Support:**

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.

---