

## COT rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A12917	Rabbit	1 mg/ml	52925

<b>Applications</b>	WB,IHC,IF,ELISA
<b>Reactivity</b>	Human,Mouse,Rat
<b>Dilution</b>	WB: 1:500 - 1:2000. IHC: 1:100 - 1:300. IF: 1:200 - 1:1000. ELISA: 1:40000. Not yet tested in other applications.
<b>Storage</b>	-20°C/1 year
<b>Specificity</b>	COT Polyclonal Antibody detects endogenous levels of COT protein.
<b>Source / Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human COT. AA range:256-305
<b>Uniprot No</b>	P41279
<b>Alternative names</b>	MAP3K8; COT; ESTF; Mitogen-activated protein kinase kinase kinase 8; Cancer Osaka thyroid oncogene; Proto-oncogene c-Cot; Serine/threonine-protein kinase cot; Tumor progression locus 2; TPL-2
<b>Form</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Conjugation</b>	
<b>Background</b>	mitogen-activated protein kinase kinase kinase 8(MAP3K8) Homo sapiens This gene is an oncogene that encodes a member of the serine/threonine protein kinase family. The encoded protein localizes to the cytoplasm and can activate both the MAP kinase and JNK kinase pathways. This protein was shown to activate I kappa B kinases, and thus induce the nuclear production of NF-kappa B. This protein was also found to promote the production of TNF-alpha and IL-2 during T lymphocyte activation. This gene may also utilize a downstream in-frame translation start codon, and thus produce an isoform containing a shorter N-terminus. The shorter isoform has been shown to display weaker transforming activity. Alternate splicing results in multiple transcript variants that encode the same protein. [provided by RefSeq, Sep 2011],
<b>Other</b>	MAP3K8, Mitogen-activated protein kinase kinase kinase 8

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation  
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster  
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All  
Species Expected

**Trademarks**

*All product names and trademarks are the property of their respective owners.*

**Regulatory Disclaimer**

*For life science research only. Not for use in diagnostic procedures.*

---

**Contact and Support:**

*To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).*

*To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.*

---