

DDR1 (Phospho-Tyr513) Antibody antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A13484	Rabbit	1 mg/ml	
Applications	WB,ELISA		
Reactivity	Human,Mouse,Rat		
Dilution	WB 1:500-2000, ELISA 1:10000-20000		
Storage	-20°C/1 year		
Specificity	This detects endogenous levels of DDR1 (Phospho-Tyr513)		
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.		
Immunogen	Synthesized phospho derived from human DDR1 (Phospho-Tyr513)		
Uniprot No	Q08345		
Alternative names	Epithelial discoidin domain-containing receptor 1 (Epithelial discoidin domain receptor 1) (EC 2.7.10.1) (CD167 antigen-like family member A) (Cell adhesion kinase) (Discoidin receptor tyrosine kinase) (HGK2) (Mammary carcinoma kinase 10) (MCK-10) (Protein-tyrosine kinase 3A) (Protein-tyrosine kinase RTK-6) (TRK E) (Tyrosine kinase DDR) (Tyrosine-protein kinase CAK) (CD antigen CD167a)		
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.		
Clonality	Polyclonal		
Isotype	IgG		
Conjugation			
Background	discoidin domain receptor tyrosine kinase 1(DDR1) Homo sapiens Receptor tyrosine kinases play a key role in the communication of cells with their microenvironment. These kinases are involved in the regulation of cell growth, differentiation and metabolism. The protein encoded by this gene belongs to a subfamily of tyrosine kinase receptors with homology to Dictyostelium discoideum protein discoidin I in their extracellular domain, and that are activated by various types of collagen. Expression of this protein is restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, it has been shown to be significantly overexpressed in several human tumors. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Feb 2011],		
Other	DDR1 CAK EDDR1 NEP NTRK4 PTK3A RTK6 TRKE, Epithelial discoidin domain-containing receptor 1 (Epithelial discoidin domain receptor 1) (EC 2.7.10.1) (CD167 antigen-like family member A) (Cell adhesion kinase) (Discoidin receptor tyrosine kinase) (HGK2) (Mammary carcinoma kinase 10) (MCK-10) (Protein-tyrosine kinase 3A) (Protein-tyrosine kinase RTK-6) (TRK E) (Tyrosine kinase DDR) (Tyrosine-protein kinase CAK) (CD antigen CD167a)		

Product Images:**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.
