

## Flg/Bek rabbit pAb antibody

| Catalog No :                 | Source:  | Concentration : | Mol.Wt. (Da): |
|------------------------------|--|-----------------|---------------|
| A14623                       | Rabbit   | 1 mg/ml         | 91868         |
| <b>Applications</b>          | WB,ELISA   |                 |               |
| <b>Reactivity</b>            | Human,Mouse,Rat  |                 |               |
| <b>Dilution</b>              | WB: 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.  |                 |               |
| <b>Storage</b>               | -20°C/1 year   |                 |               |
| <b>Specificity</b>           | Flg/Bek Polyclonal Antibody detects endogenous levels of Flg/Bek protein.  |                 |               |
| <b>Source / Purification</b> | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |                 |               |
| <b>Immunogen</b>             | Synthesized peptide derived from Flg/Bek . at AA range: 410-490  |                 |               |
| <b>Uniprot No</b>            | P11362/P21802  |                 |               |
| <b>Alternative names</b>     | FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331; FGFR2; BE   |                 |               |
| <b>Form</b>                  | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |                 |               |
| <b>Clonality</b>             | Polyclonal   |                 |               |
| <b>Isotype</b>               | IgG  |                 |               |
| <b>Conjugation</b>           |  |                 |               |
| <b>Background</b>            | <p>fibroblast growth factor receptor 1(FGFR1) Homo sapiens The protein encoded by this gene is a member of the fibroblast growth factor receptor (FGFR) family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member binds both acidic and basic fibroblast growth factors and is involved in limb induction. Mutations in this gene have been associated with Pfeiffer syndrome, Jackson-Weiss syndrome,</p> |                 |               |
| <b>Other</b>                 | FGFR1/FGFR2, Basic fibroblast growth factor receptor 1/2   |                 |               |

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

**Trademarks**

All product names and trademarks are the property of their respective owners.

**Regulatory Disclaimer**

For life science research only. Not for use in diagnostic procedures.

---

**Contact and Support:**

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.

---