

## FOG1 rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A14673	Rabbit	1 mg/ml	
<b>Applications</b>	WB,ELISA		
<b>Reactivity</b>	Human,Mouse		
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000		
<b>Storage</b>	-20°C/1 year		
<b>Specificity</b>	FOG1 Polyclonal Antibody detects endogenous levels of protein.		
<b>Source / Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.		
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 620-700		
<b>Uniprot No</b>	Q8IX07		
<b>Alternative names</b>			
<b>Form</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.		
<b>Clonality</b>	Polyclonal		
<b>Isotype</b>	IgG		
<b>Conjugation</b>			
<b>Background</b>	<p>domain:The CCHC-type zinc fingers 1, 5, 6 and 9 directly bind to GATA-type zinc fingers. The Tyr residue adjacent to the last Cys of the CCHC-type zinc finger is essential for the interaction with GATA-type zinc fingers.,function:Transcription regulator that plays an essential role in erythroid and megakaryocytic cell differentiation. Essential cofactor that acts via the formation of a heterodimer with transcription factors of the GATA family GATA1, GATA2 and GATA3. Such heterodimer can both activate or repress transcriptional activity, depending on the cell and promoter context. The heterodimer formed with GATA proteins is essential to activate expression of genes such as NFE2, ITGA2B, alpha- and beta-globin, while it represses expression of KLF1. May be involved in regulation of some genes in gonads. May also be involved in cardiac development, in a non-redundant way with ZFPM2/FOG2.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the FOG (Friend of GATA) family.,similarity:Contains 4 C2H2-type zinc fingers.,similarity:Contains 5 C2HC-type zinc fingers.,subunit:Interacts with corepressor CTBP2; this interaction is however not essential for corepressor activity (By similarity). Interacts with the N-terminal zinc-finger of GATA1, GATA2 and probably GATA3.,tissue specificity:Mainly expressed in hematopoietic tissues. Also expressed in adult cerebellum, stomach, lymph node, liver and pancreas. Expressed in fetal heart, liver and spleen..</p>		

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry CHIP-Chromatin Immunoprecipitation  
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster  
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All  
Species Expected

**Trademarks**

*All product names and trademarks are the property of their respective owners.*

**Regulatory Disclaimer**

*For life science research only. Not for use in diagnostic procedures.*

---

**Contact and Support:**

*To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).*

*To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.*

---