

## Histone H4 (Tri Methyl Lys20) rabbit pAb antibody

| Catalog No :                 | Source:  | Concentration : | Mol.Wt. (Da): |
|------------------------------|--|-----------------|---------------|
| A15760                       | Rabbit   | 1 mg/ml         | 11367         |
| <b>Applications</b>          | WB   |                 |               |
| <b>Reactivity</b>            | Human,Mouse,Rat  |                 |               |
| <b>Dilution</b>              | WB: 1:500-1000   |                 |               |
| <b>Storage</b>               | -20°C/1 year   |                 |               |
| <b>Specificity</b>           | The antibody detects endogenous Histone H4 (Tri Methyl Lys20) protein.   |                 |               |
| <b>Source / Purification</b> | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using specific immunogen.  |                 |               |
| <b>Immunogen</b>             | Synthetic Peptide of Histone H4 (Tri Methyl Lys20)   |                 |               |
| <b>Uniprot No</b>            | P62805   |                 |               |
| <b>Alternative names</b>     | H4K20ME3; HIST1H4A; H4/A; H4FA; HIST1H4B; H4/I; H4FI; HIST1H4C; H4/G; H4FG; HIST1H4D; H4/B; H4FB; HIST1H4E; H4/J; H4FJ; HIST1H4F; H4/C; H4FC; HIST1H4H; H4/H; H4FH; HIST1H4I; H4/M; H4FM; HIST1H4J; H4/E; H4FE; HIST1H4K; H4/D; H4FD; HIST1H4L; H4/K; H4FK; HIST2H4A; H4/N; H4F2; H4FN; HIST2H4; HIST2H4B; H4/O; H4FO; HIST4H4; Histone H4   |                 |               |
| <b>Form</b>                  | PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.  |                 |               |
| <b>Clonality</b>             | Polyclonal   |                 |               |
| <b>Isotype</b>               | IgG  |                 |               |
| <b>Conjugation</b>           |  |                 |               |
| <b>Background</b>            | histone cluster 1 H4 family member i(HIST1H4I) Homo sapiens Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H4 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the histone microcluster on chromosome 6p21.33. [provided by RefSeq, Aug 2015], |                 |               |
| <b>Other</b>                 | HIST1H4A, Histone H4   |                 |               |

---

**Product Images:****Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

**Species Cross-Reactivity Key:**

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

**Trademarks**

All product names and trademarks are the property of their respective owners.

**Regulatory Disclaimer**

For life science research only. Not for use in diagnostic procedures.

---

**Contact and Support:**

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit [www.aabsci.cn](http://www.aabsci.cn), contact information will be displayed.

---