

VSV-G-Tag mouse mAb(8D6) antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (Da):
A42057	Mouse	1 mg/ml	

Applications	WB,IP,IF
Reactivity	Species independent
Dilution	WB: 1:5000 ; IP: 1:200; IF: 1:1000
Storage	-20°C/1 year
Specificity	The antibody detects C-terminal, internal, and N-terminal VSV-G fusion proteins.
Source / Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Immunogen	Synthetic Peptide of VSV-G-Tag
Uniprot No	
Alternative names	
Form	PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.
Clonality	Monoclonal
Isotype	
Conjugation	
Background	The fusiogenic envelope G glycoprotein of the vesicular stomatitis virus (VSV-G) that has been used to pseudotype retrovirus and lentivirus vectors can be used alone as an efficient vehicle for gene transfer. The VSV-G epitope tag is commonly engineered onto the N- or C- terminus of a protein of interest so that the tagged protein can be analyzed and visualized using immunochemical methods.

Other

Product Images:

Application Key:

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.