

MED28 rabbit pAb antibody

Applications :	Source:	Concentration :	Mol.Wt. (Da):
Reactivity A1/404	Human, Mouse, Rat Rabbit	1 mg/ml	
Dilution	WB 1:500-2000 ELISA 1:5000-20000		
Storage	-20°C/1 year		
Specificity	MED28 Polyclonal Antibody detects endogenous levels of protein.		
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.		
Immunogen	Synthesized peptide derived from human protein . at AA range: 20-100		
Uniprot No	Q9H204		
Alternative names			
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.		
Clonality	Polyclonal		
Isotype	IgG		
Conjugation			
Background	<p>function:Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. May be part of a complex containing NF2/merlin that participates in cellular signaling to the actin cytoskeleton downstream of tyrosine kinase signaling pathways.,induction:Up-regulated by endothelial cells when exposed to tumor conditional media.,similarity:Belongs to the Mediator complex subunit 28 family.,subcellular location:According to PubMed:15467741, it is cytoplasmic and mainly membrane-associated.,subunit:Component of the Mediator complex, which is composed of MED1, MED4, MED6, MED7, MED8, MED9, MED10, MED11, MED12, MED13, MED13L, MED14, MED15, MED16, MED17, MED18, MED19, MED20, MED21, MED22, MED23, MED24, MED25, MED26, MED27, MED29, MED30, MED31, CCNC, CDK8 and CDC2L6/CDK11. The MED12, MED13, CCNC and CDK8 subunits form a distinct module termed the CDK8 module. Mediator containing the CDK8 module is less active than Mediator lacking this module in supporting transcriptional activation. Individual preparations of the Mediator complex lacking one or more distinct subunits have been variously termed ARC, CRSP, DRIP, PC2, SMCC and TRAP. Forms a ternary complex with NF2/merlin and GRB2. Binds to actin.,tissue specificity:Widely expressed. Highly expressed in vascular tissues such as placenta, testis and liver.,</p>		

Product Images:**Application Key:**

WB-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation

IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster

X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All

Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.cn, contact information will be displayed.
